



Short Communication

Saissetia coffeae (Walker) (Coccidae: Hemiptera) infestation on *Meyna spinosa* Roxb. ex Link-a new host plant and sap-sucker (pest) record from Assam, India

Ankur Jyoti Saikia, Vipin Parkash, Kalyan Das, Pallwabee Duarah

Abstract

The hemispherical scale insect *Saissetia coffeae* (Walker) is reported for the first time from Assam, India infesting one-year old *Meyna spinosa* Roxb. ex Link (Rubiaceae) seedlings in nursery conditions. The host plant is attributed with medicinal properties amongst the inhabitants of the region; although occurring mostly as scanty populations in the wild, probably a cause of pest infestations.

Keywords medicinal plants pests, new record; polyphagy, sap sucker, scale insect

Introduction

Scale insects or Coccoids are sap-sucking hemipterans characterized as paedomorphic hemipterans and completely metamorphosed males [1]. The existence of protective covering/ scale and a small, cryptic habit are amongst the main features of this group of insects [2]. Scale insects, basically suck the plant sap which affects its growth, leads to defoliation, withering of shoots/entire plant. A secretion of honeydew which coats the plant surface impedes the assimilation and photosynthesis and it is also a perfect medium for sooty mould [3-5].

Saissetia coffeae (Walker), commonly known as hemispherical scale insect is a polyphagous insect and a pest of several vegetable and fruit crops [6]. Its main damage is the secretion of profuse amounts of honeydew; later colonized by black sooty mold, contaminating foliar abaxial surfaces. The honeydew is also observed to attract ants which might protect the pest from its natural enemies.

Meyna spinosa Roxb. ex Link (Rubiaceae), also known as *Kutkura* (Assamese) and *Lam heibi* (Manipuri) is a popular *albeit* wild plant of ethnomedicinal usage in hepatic, gastrointestinal and dermatic ailments amongst the people North-Eastern regions of India [7].

Methodology

In course of raising *M. spinosa* seedlings, the authors observed aforementioned pests on one-year old seedlings raised in the nursery at Deovan campus of Rain Forest Research Institute, Sotai, Jorhat, Assam during April-June, 2016 (Figure 1. A-D). Specimens were collected in 70% ethanol and were slide mounted as per the method of Hodgson and Henderson [8]. Literature survey [8] helped in the identification of the collected scale insect. Photographs were taken using a CANON power Shoot G/0/11 (digital) camera under a Stereozoom

Received: 21 December 2017

Accepted: 22 February 2018

Online: 28 February 2018

Authors:

A. J. Saikia ✉

Soil Microbiology Laboratory - I, Silviculture & Forest Management Division, Rain Forest Research Institute (Indian Council of Forestry Research and Education, An Autonomous Council under Ministry of Environment, Forests and Climate Change, Government of India), Deovan, Jorhat-785001, Assam, India

V. Parkash

Forest Research Institute (Deemed) University (Indian Council of Forestry Research & Education) An Autonomous Council under Ministry of Environment, Forests and Climate Change, Government of India), Dehradun-248195, (Uttarakhand) India

K. Das

Department of Botany, Jagannath Barooah College (Autonomous), Jorhat, Assam, India

P. Duarah

Department of Zoology, Jagannath Barooah College (Autonomous), Jorhat, Assam, India

✉ ankurjyoti05@rediffmail.com

Emer Life Sci Res (2018) 4(1): 15-18

E-ISSN: 2395-6658

P-ISSN: 2395-664X

DOI: <https://doi.org/10.31783/elsr.2018.411518>

Trinocular Microscope (Make: LABOVISION, Model: AZM 100)



Figure 1. (A-D): *Saissetia coffeae* infestation on stem, spines and foliage of *Meyna spinosa*;
(E) Dorsal view of adult *S. coffeae*; (F) Dorsal view of *S. coffeae* nymph (with the characteristic H mark)



Results and Discussion

Saissetia coffeae (Walker)

Synonyms: *Lecanium coffeae* Walker, 1852: 1079; *Lecanium hemisphaericum* Targioni Tozzetti, 1867: 26; Cockerell & Parrott, 1899: 164; *Chermes filicum* Boisduval, 1867: 328; *Saissetia filicum* (Boisduval); *Chermes hibernaculorum* Boisduval, 1867: 328; *Lecanium coffeae* Walker, 1852: 1079; *Lecanium hemisphaericum* Targioni Tozzetti, 1867: 26; Cockerell & Parrott, 1899: 164; *Chermes filicum* Boisduval, 1867: 328; *Saissetia filicum* (Boisduval); –Fernald, 1903: 201; *Chermes hibernaculorum* Boisduval, 1867: 328.

Material examined

Meyna spinosa Roxb. ex Link (Rubiaceae) seedling leaves, petiole and branches, Deovan campus, Rain Forest Research Institute, Sotai, Jorhat, Assam. Coll. A. J. Saikia (Figure 1. A-D).

Morphological Diagnosis: (Figure 1. E-F)

Unmounted material

Adult-convex and rounded; color shiny tan; dorsal surface completely smooth; 2 mm long; Nymph-yellowish-green in colour, characteristic ‘H’ mark on dorsal surface

Mounted material

Dorsum-heavily sclerotised, 4µm long dorsal setae frequent throughout dorsum, **Margin**-Setae of 2 sizes, longer between stigmatic areas and shorter between anterior stigmatic clefts, **Venter**-well developed legs, claws without a denticle.

Distribution range

The scale insect is found throughout the tropics as well as in some other sub-tropical areas as well [9]. Its distribution range encompasses Asia, Africa, North America, South America, Europe and Oceania [10].

Damage potential

It is a pest to cultivated plants such as guava, coffee, cotton, eggplant, okra, citrus, mango, tea, banana, etc. It also infests wildy occurring plants [9].

Notes

(I) *Meyna spinosa*, is attributed with ethnomedicinal claims by the inhabitants of this region. Although producing a good seed set and exhibiting the above average seed germination percentage; it is found in scarcity. The present study investigations have shown the deformation and wilting of the plants infested by the *S. coffeae* pest. It has also been reported that the “black looper” caterpillar (*Hyposidra talaca* Walker) occurs as a pest for this plant. Synergistically, these two pests can be attributed to lowering the occurrence of the plant; despite of a good seed set [11].

(II) The present inspection has also re-established the polyphagous nature of the pest and added an extension of a new host in the established lists [6, 8]. Although occurring in the geographical entity (state/country), it has no record of the observed plant species as a host [10]

Conclusion

The family Coccidae is an important group because of its role as agricultural pests or biocontrol agent and/or pests in many ecosystems. *Meyna spinosa* Roxb. ex Link is a new host recorded for *Saissetia coffeae* (Walker) in the northeastern part of the Indian sub-continent, a confluence of two biodiversity hotspots, viz.- the Indo-Myanmar and the Eastern Himalayas [12] in this study.



Acknowledgements

Authors like to thank Shri Pabitra Saikia, multi-tasking staff and Sri Prodip Bora, Semi-skilled laborer who assisted in collection of *M. spinosa* seeds and raising of seedlings. Authors AJS and VP would like to thank the Director, Rain Forest Research Institute, Jorhat, Assam (India) for laboratory facilities; while KD and PD would like to extend their gratitude to the Principal, J. B. College (Autonomous), Jorhat, Assam, India

Transparency declaration

The authors declare no conflicts of interest.

References

- [1] P. J. Gullan and J. H. Martin (2003). Sternorrhyncha (jumping plant-lice, white flies, aphids and scale insects). In: Resh, V. and Cardé, R. Encyclopedia of Insects, Academic Press Amsterdam, pp. 1079-1089.
- [2] P. J. Gullan and L. G. Cook (2007). Phylogeny and higher classification of scale insects (Hemiptera: Sternorrhyncha:Coccoidea). Zootaxa, **1668**: 413-425
- [3] A. Dziedzicka (1988). Welnowce szklarniowe (Homoptera, Coccinea, Pseudococcidae) [in Polish]. Zeszyty Problemowe Postępów Nauk Rolniczych, **333**: 87-91.
- [4] M. Kosztarab (1996). Scale insects of Northeastern North America. Virginia Museum of Natural History, Martinsville, pp. viii – 650.
- [5] Y. Ben-Dov and C. J. Hodgson (1997). Soft scale insects-Their Biology, Natural Enemies and Control. Elsevier, Amsterdam and New York, pp. 439.
- [6] R. C. Henderson, A. Sultan and A. W. Robertson (2010). Scale insect fauna (Hemiptera: Sternorrhyncha: Coccoidea) of New Zealand's pygmy mistletoes (*Korthalsella*: Viscaceae) with description of three new species: *Leucapsis albotecta*, *L. trilobata* (Diaspididae) and *Eriococcus korthalsellae* (Eriococcidae). Zootaxa, **2644**: 1-24.
- [7] N. K. S. Singh, W. Radhapiyari, Md. A. H. Choudhury, Th. B. Singh, Kh. Bonny and Th. Sobita Devi (2015). Spectroscopic study of antioxidant property and trace elements of *Meyna spinosa* Roxb. ex Link leaves. Indian Journal of Natural Products and Resources, **6**: 51-55.
- [8] C. J. Hodgson and R. C. Henderson (2000). Coccidae (Insecta: Hemiptera: Coccoidea). Fauna of New Zealand, 41.
- [9] D. Nafus (2000). Hemispherical scale Insect (*Saissetia coffeae* [Walker]) Agricultural Pests of the Pacific. ADAP, Honolulu, US. https://www.ctahr.hawaii.edu/adap/Publications/ADAP_pubs/2000-8.pdf (accessed on 19 July, 2016).
- [10] CABI (2017). *Saissetia coffeae* (hemispherical scale). In: Invasive Species Compendium. Wallingford, UK: CAB International. <http://www.cabi.org/isc/datasheet/48202> (accessed on 19 July, 2016).
- [11] A. J. Saikia and V. Parkash (2017). *Hyposidra talaca* Walker (Geometridae: Lepidoptera) a New Pest (Defoliator) Recorded for *Meyna spinosa* Roxb. ex Link from Assam, India. Indian Forester, **143**: 714-715.
- [12] Conservation International (2004). Hot Spots Revisited. <https://biodiversityhotspots.org/XP/hotspots/himalaya> (accessed on 19 July, 2016).